

MicroRNAs in Atrial Fibrillation: Pathophysiological Aspects and Potential Biomarkers

Aleksey M. Chaulin, MD^{1,2*}; Dmitriy V. Duplyakov, PhD, ScD^{1,2}

¹Samara State Medical University

²Samara Regional Clinical Cardiology Dispensary
 Samara, the Russian Federation

Abstract

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia associated with increased morbidity and mortality. The pathophysiological mechanisms underlying the development of AF are not entirely clear. In addition, there are no optimal biomarkers for the early detection and prognosis for patients with AF. Emerging studies have uncovered a role for miRNAs in the initiation and maintenance of cardiovascular disease. This review discusses the role of miRNAs in the pathogenesis of AF as well as the possibility of using miRNAs as biomarkers for detecting and predicting AF. (**International Journal of Biomedicine. 2020;10(3):198-205.**)

Key Words: atrial fibrillation • microRNA • pathophysiology • biomarker

Abbreviations

AF, atrial fibrillation; ANS, autonomic nervous system; CAF, chronic AF; CHD, coronary heart disease; CVD, cardiovascular disease; CHF, congestive heart failure; ER, electrical remodeling; HF, heart failure; LAA, left atrial appendage; LTCC, L-type calcium channel; LA, left atria; miRNA, microRNA; c-miRNA, circulating miRNA; RAA, right atrial appendage; SR, structural remodeling.

Atrial fibrillation (AF) is the most common sustained cardiac arrhythmia. According to the latest data, there are more than 33 million people suffering from AF in the world, and gender-specific morbidity has been observed. In men, the incidence of AF is 3 times higher than in women.⁽¹⁾ Many genetic, molecular, and environmental risk factors associated with AF are also major risk factors for morbidity and mortality among patients with cardiovascular disease (CVD).⁽²⁾

The simplest and most frequently used method for diagnosing AF is electrocardiography, but this method has a significant drawback, which consists in the short duration of recording the electrical activity of the heart, which limits the diagnosis in asymptomatic patients.⁽³⁾ Existing laboratory biomarkers of myocardial damage (cardiac troponins, natriuretic peptides, creatine phosphokinase, lactate dehydrogenase, aspartate aminotransferase, etc.), among which cardiac troponins

are most sensitive and specific,^(4,5) are not very effective for detecting and predicting AF.⁽⁶⁾ The elevation in troponin levels occurs because of a mismatch between oxygen demand and supply. The escalated heart rate increases myocardial oxygen demand while decreasing myocardial oxygen supply by shortening diastolic time.^(6,7) The shortened diastolic phase reduces the time for myocardial perfusion, a majority of which occurs during diastole. According to some reports, troponins correlate well with the severity of arrhythmias and can be used as prognostic markers,⁽⁸⁾ but according to other sources, they are ineffective as biomarkers.⁽⁹⁾ Therefore, there is a need to search for new and reliable biomarkers for the early diagnosis of AF.

Heart failure (HF), diabetes, hypertension, hyperthyroidism, obesity, gender, and structural and ischemic heart diseases are the most common risk factors associated with AF.⁽¹⁰⁾ The genetic component was recently also considered as a significant risk factor for AF.^(11,12)

miRNAs are endogenous 21–22-nucleotide noncoding small RNAs that regulate physiological and disease processes at the post-transcriptional level. The first such tiny regulatory RNA

*Corresponding author: Aleksey M. Chaulin, MD.
 Samara State Medical University; Samara Russia. E-mail:
alekseymichailovich22976@gmail.com

to be identified was the lin-4 RNA, which controls the timing of *Caenorhabditis elegans* larval development.⁽¹³⁾ Over the last few years, miRNAs have been the focus of many studies, providing evidence that these small RNAs have a significant role in the modification of numerous biological pathways.⁽¹⁴⁾ miRNAs regulate approximately 30% of all protein-coding genes and have therefore been predicted to be involved in almost all cellular processes.⁽¹⁵⁾ miRNAs have been implicated in various human diseases like cancer,⁽¹⁶⁻¹⁸⁾ CVD,⁽¹⁹⁻²¹⁾ neurological diseases,^(22,23) autoimmune diseases,⁽²⁴⁻²⁶⁾ and asthma.^(27,28)

miRNA regulation in CVD was first described in 2006, when specific miRNAs were found to be up- or downregulated in mouse models of cardiac hypertrophy and HF.⁽²⁹⁾ Later, it was demonstrated that AF is associated with altered miRNA levels in atrial tissue and plasma.^(30,31) The regulatory function of specific miRNAs has been studied in the structural and electrical remodeling of the heart.⁽³²⁻³⁶⁾ A number of studies have shown that c-miRNAs associated with AF could serve as potential biomarkers of the disease, whereas specific tissue miRNAs could become targets for therapy.⁽³⁷⁻³⁸⁾

According to the latest data, miRNAs play a leading role in the pathophysiology of AF, regulating mechanisms of atrial remodeling. The reduced or increased expression of miRNAs is genetically programmed and under normal conditions contributes to the development of the cardiovascular system in humans and animals.⁽³⁹⁾ At the same time, a change in the level of miRNA expression in circulating blood and tissues can occur under pathological conditions associated with the development of various CVDs, including AF, which lead to myocardial remodeling. Numerous experimental and clinical studies found the role that miRNAs play in the pathogenesis of AF. The following main pathogenetic links in the initiation and progression of AF can be distinguished, in the regulation of which microRNA is involved: electrical remodeling (ER), structural remodeling (SR), remodeling of ANS, and dysregulation of calcium levels. A discussion of the involvement of miRNAs in each of these mechanisms is presented in the main part of this article.

Electrical remodeling in AF mediated by miRNAs

ER is the most common change associated with AF. ER occurs due to decreasing L-type calcium channel conductance (ICaL) and increasing inwardly rectifying potassium conductance (IK1). Changes in the electrical properties of Ca2+-dependent potassium channels and *connexin* channels also cause AF-related ER.⁽⁴⁰⁾ In the process of ER, several miRNAs are involved: miRNA-1, miRNA-328 and miRNA-499, each of which exhibits several specific properties.

miRNA-1

miRNA-1 is abundantly expressed in the myocardium and plays an important role in the development of the heart and electrical activity. Dysregulation in miRNA-1 expression leads to myocardial hypertrophy, myocyte proliferation, and cardiac arrhythmias. In addition, the arrhythmogenic potential of miRNA-1 is associated with CHD.⁽⁴¹⁾ A study on a rabbit model of atrial tachycardia showed that overexpression of miRNA-1 shortens the atrial refractory period induced by tachycardia and increases inward-rectifier current activity (IKs) by reducing the expression of the *KCNE1* and *KCNB2*

genes encoding the potassium channel subunits. miRNA-1 knockdown, in contrast, weakened the suppression of the *KCNE1* and *KCNB2* genes, resulting in a shortening of the atrial effective period and an increase in IKs. Thus, it was found that *KCNE1* and *KCNB2* are target genes for miRNA-1. It has also been suggested that targeted suppression of these potassium channel genes enhances the duration and frequency of AF.

Jia et al.⁽⁴²⁾ showed the critical role of miRNA-1 in atrial ER and the clinical importance of miRNA-1 as a therapeutic agent in AF. Tsoporis et al.⁽⁴³⁾ studied the expression of the muscle-specific miRNA-1 and miRNA-133A in RAA biopsies and blood plasma taken before aortic cross-clamping and after reperfusion. There was no change in tissue or plasma miRNA-1 and miRNA-133A levels compared to pre CABG. However, in patients who postoperatively developed AF, compared to patients who remained in sinus rhythm, tissue miRNA-1 increased whereas miRNA-133A decreased and negatively correlated with RAA apoptosis. In postoperative AF, differential regulation of pro- and anti-apoptotic miRNAs-1 and -133A respectively in the RAA, may contribute to postoperative apoptosis. These results provide new insights into molecular mechanisms of postoperative AF with potential therapeutic implications.

As is known, miRNA-1 and miRNA-133 regulate the expression of multiple genes involved in myocardial function, including hyperpolarization-activated cyclic nucleotide-gated (HCN) cation channels. In the study performed by Yao-Dong Li et al.,⁽⁴⁴⁾ aged patients with AF, compared with aged patients with sinus rhythm, exhibited significantly higher HCN2 and HCN4 channel mRNA and protein expression levels ($P < 0.05$), but significantly lower expression levels of miRNA-1 and miRNA-133 ($P < 0.05$). Expression levels of HCN2 and HCN4 increased with age, and a greater increase was identified in patients with age-associated atrial fibrillation compared with that in those with aged sinus rhythm. These electrophysiological changes may contribute to the induction of ectopic premature beats that trigger AF. In patients with AF, miRNA-1 levels are greatly reduced, possibly contributing to up-regulation of Kir2.1 subunits, leading to increased IK1. Because up-regulation of inward-rectifier currents is important for AF maintenance, these results provide potential new insights into molecular mechanisms of AF with potential therapeutic implications.⁽⁴⁵⁾

Numerous studies have confirmed the relationship between the changes in expression profile of c-miRNAs and the regulatory effect of AF-related miRNAs on ion channels. The results obtained by Yingmin Lu et al.⁽⁴⁶⁾ indicated that transfected miRNA-1 could significantly inhibit the expression of CACNB2 protein. miRNA-1 can decrease the intracellular Ca(2+) concentration and prevent the AF. In addition, the recently proposed involvement of miRNA-1 in the modulation of Ca2+ handling proteins, including calmodulin, phospholamban, Na+/Ca2+ exchanger (NCX), sorcin, junctin, triadin, eventually resulting in shortened refractoriness of sarcoplasmic reticulum Ca2+ release further support the hypothesis of a pivotal functional role of miRNA-1 in the pathogenesis of AF.⁽⁴⁷⁾

miRNA-328

miRNA-328 contributes to adverse atrial ER in AF through targeting L-type Ca(2+) channel genes. A potential role

for miRNA regulation of cardiac depolarization was identified when Lu et al.⁽⁴⁸⁾ utilized computational prediction algorithms to identify *CACNA1C* and *CACNB1* as potential targets for miRNA-328. Overexpression of miRNA-328 in experimental models recapitulated the phenotypes of AF, exemplified by enhanced AF vulnerability, diminished L-type Ca^{2+} current, and shortened potential duration of atrial action. *CACNA1C* and *CACNB1* were established as the cognate target genes for miRNA-328 by western blot and luciferase activity assay, showing the reciprocal relationship between the levels of miRNA-328 and I CaL channel protein subunits.^(48,49) Soeki et al.⁽⁵⁰⁾ found that plasma levels of miRNA-328 were higher in AF patients than in control subjects. Plasma miRNA-328 levels were significantly higher in the LAA than in the periphery and PV in patients with AF, but not in control subjects. The authors concluded that local production of miRNA-328 in the LA may be involved in the process of atrial remodeling in AF patients.

miRNA-499

Recently, *KCNV3*, the gene that encodes the small-conductance, calcium-activated potassium channel 3 (SK3), was found to be strongly associated with AF. The study performed by Ling et al. showed that in patients with permanent AF, atrial miRNA-499 was significantly upregulated, leading to SK3 downregulation through effects on the *KCNV3* gene expression and possibly contributing to ER in AF.⁽⁵¹⁾ In addition, upregulation of atrial miRNA-499 induced the downregulation of *CACNB2* (an important subunit of the LTCC) expression and may contribute to ER in AF.⁽⁵²⁾ Increased expression of miRNA-499 (2.3-fold) was observed in patients with acute new-onset AF, compared with well-controlled AF and control patients.⁽⁵³⁾

Structural remodeling in AF mediated by miRNAs

miRNAs that participate in sarcoplasmic reticulum regulate the genes encoding the proteins responsible for the formation of the extracellular matrix and contribute to atrial fibrosis by regulating pro- and antifibrotic signaling pathways.⁽⁵⁴⁾ These miRNAs are mainly involved in a decrease in the conduction rate and an increase in the reentrant activity.⁽⁵⁵⁾ The following miRNAs participate in SR of the heart in AF: miRNA-21, miRNA-29, miRNA-126, miRNA-150, and miRNA-483.

miRNA-21

Barana et al.⁽⁵⁶⁾ reported that chronic AF (CAF) increases miRNA-21 expression in enzymatically isolated human atrial myocytes. Moreover, it decreases I CaL density by downregulating the expression of Ca^{2+} channel subunits (*CACNA1C* and *CACNB2*). These results suggest that miRNA-21 could participate in the CAF-induced I CaL downregulation and in shortening the duration of the action potential that maintains the arrhythmia.

Atrial fibrosis is the most important substrate of AF. In a study performed by Adam et al.,⁽⁵⁷⁾ LA from patients with AF showed a 2.5-fold increased expression of miRNA-21, compared to matched LA of patients in sinus rhythm. Increased miRNA-21 expression correlated positively with atrial collagen content and was associated with a reduced protein expression of Spry1 and increased expression of connective tissue growth factor, lysyl oxidase and Rac1-GTPase.⁽⁵⁷⁾

Thum et al.⁽⁵⁸⁾ showed that miRNA-21 regulates the

ERK-MAP kinase signaling pathway in cardiac fibroblasts, which impacts global cardiac structure and function. In vivo silencing of miRNA-21 by a specific antagomir in a mouse pressure-overload-induced disease model reduces cardiac ERK-MAP kinase activity, inhibits interstitial fibrosis and attenuates cardiac dysfunction.

The PTEN/PI3K signaling pathway was involved in a variety of physiological activities, including cell proliferation, differentiation, and apoptosis, which participate in the pathophysiology of various diseases.⁽⁵⁹⁾ The target genes of miRNA-21, namely, *PTEN*, are PI3K signaling pathway inhibitors. Overexpression of miRNA-21 decreased expressions of *PTEN*, thereby activating the PI3K signaling pathway, which may play a crucial role in initiation and maintenance of atrial SR.⁽⁶⁰⁾

miRNA-29

The human miRNA-29 family of mirnas has three mature members, miRNA-29a, miRNA-29b, and miRNA-29c. miRNA-29s directly target at least genes related to the extracellular matrix. These genes code for several of the key proteins involved in the physiological or pathological formation of extracellular matrix. The regulation of the extracellular matrix by miRNA-29s has been implicated in the development of fibrosis in many organs, including the heart.⁽⁶¹⁾ Strong antifibrotic effects of miRNA-29s have been demonstrated in the heart, kidney, and other organs.⁽⁶²⁾ Expression of miRNA-29b extracellular matrix target genes collagen-1A1 (*COL1A1*), collagen-3A1 (*COL3A1*), and fibrillin increased significantly in CHF fibroblasts. miRNA-29b plasma levels were decreased in patients with CHF or AF (by 53% and 54%, respectively; both $P < 0.001$) and were further decreased in patients with both AF and CHF (by 84%; $P < 0.001$). miRNA-29b expression was also reduced in the atria of CAF patients (by 54% versus sinus rhythm; $P < 0.05$). Adeno-associated virus-mediated knockdown of miRNA-29b in mice significantly increased atrial *COL1A1* mRNA expression and cardiac tissue collagen content. miRNA-29 likely plays a role in atrial fibrotic remodeling and may have value as a biomarker and/or therapeutic target.⁽⁶³⁾

miRNA-126

miRNA-126 plays an active role in angiogenesis by regulating the expression of Epidermal Growth Factor Like-domain 7 (EGFL7). EGFL7 has been identified as a potential miRNA-126 target. EGFL7 and its specific miRNA-126 may be involved in the pathogenesis of vasculopathy and fibrosis.⁽⁶⁴⁾ In a study by Wei et al.,⁽⁶⁵⁾ relative miRNA-126 expression was downregulated in patients with AF and/or HF compared with controls. Moreover, miRNA-126 expression was markedly lower in the HF-AF group compared with the AF and HF groups. Multiple linear regression analysis showed that miRNA-126 expression was positively correlated with LVEF, but negatively correlated with the logarithm of NT-pro BNP and the cardiothoracic ratio (all $P < 0.05$). Thus, serum miRNA-126 levels could serve as a potential candidate biomarker for evaluating the severity of AF and HF.

miRNA-150

In a study performed by McManus,⁽⁶⁶⁾ plasma levels of miRNA-150, known to influence atrial fibrosis and pro-fibrillatory potential under experimental conditions, were

2-fold lower among patients with AF than among those without AF. In addition, plasma level of miRNA-150 also was lower in participants with persistent AF than in those with paroxysmal AF and increased 3-fold after AF ablation. Goren et al.⁽⁶⁷⁾ found that miRNA-150 expression was 3.2-fold lower in platelets of HF patients with AF, relative to those without AF. A similar effect was seen in serum samples from the same patients, in which miRNA-150 levels were 1.5-fold lower ($P=0.004$) in HF patients with AF. Thus, the authors concluded that miRNA-150 expression levels in platelets of patients with systolic HF with AF are significantly reduced and correlated to the cell-free circulating levels of this miRNA.

miRNA-483

miRNA-483-5p, typically transcribed with its host gene, *IGF2*, has been isolated in several human tissue types including myocardium, hepatic and brain tissue as well as in blood serum.^(68,69) Harling et al.⁽⁷⁰⁾ found significant overexpression of miRNA-483-5p in the pre-operative serum of patients with postoperative AF. It is possible that the observed increase in miRNA-483-5p expression represents upregulation of the *IGF2* gene, which, for example, may be stimulated in states of cardiac stress. Overexpression of *IGF2* may also play a role in providing a pro-inflammatory substrate through the regulation of NF- κ B- and IL-6-mediated pathways, which may in turn lower the threshold on which a surgical trigger may potentiate AF in the postoperative period.⁽⁷¹⁾

Remodeling of the autonomic nervous system in AF mediated by miRNAs

ANS has been shown to have an important role in the initiation and maintenance of AF.⁽⁷²⁾ Activation of either the extrinsic parasympathetic or sympathetic neural elements of the cardiac ANS has been reported to induce rapid focal firing and AF induction via mechanisms of shortening the atrial or pulmonary vein refractoriness mediated by parasympathetic neurotransmitters or increasing intracellular Ca^{2+} concentrations mediated by sympathetic neurotransmitters, respectively. These results indicate the complexity and multiple functions of the cardiac ANS in inducing AF.⁽⁷³⁾

Rao et al.,⁽⁷⁴⁾ utilizing the right cervical vagus trunk stimulation (RVTS) to successfully induce AF by reducing the effective refractory period (ERP) and increasing dERP, hypothesized that RVTS activates the extrinsic and/or intrinsic cardiac ANS to mediate neural remodeling, which initiates and/or maintains AF, presumably by increasing the activity of acetylcholine. Modulation of ANS activity may constitute an important therapeutic strategy for the management of atrial tachyarrhythmias.⁽⁷⁵⁾

At least two miRNAs, miRNA-30 and miRNA-206, are involved in the ANS remodeling in AF.

miRNA-30

Among 355 of 1,205 human miRNAs being expressed and assumed functioning in the heart, miRNA-30d is the sixth most abundantly expressed miRNA, which indicates that it could be a crucial regulator of cardiomyocytes in the physiological and/or pathophysiological condition. Morishima et al.⁽⁷⁶⁾ demonstrated a novel role for miRNA-30d in remodeling the cardiac ion channel in human persistent AF. miRNA-30d was upregulated, whereas *KCNJ3*, a target gene of miRNA-30d,

was downregulated in human AF cardiomyocytes. The authors provided the first evidence that the downward remodeling of IK.ACh in Ca^{2+} -overloaded cardiomyocytes is attributed, at least in part, to deranged Ca^{2+} handling, leading to the upregulation of miRNA-30d in human AF.

miRNA-206

Cardiac autonomic nerve remodeling (ANR) is a critical mechanism in AF. Zhang et al.⁽⁷⁷⁾ performed a study to assess the underlying role of miRNAs in regulating cardiac ANR in AF by right atrial tachypacing (A-TP) in dogs. Forced expression of miRNA-206 through lentiviral infection based on A-TP *in vivo* significantly shortened the atrial ERP. The regeneration of nerves increased more than 2-fold by miRNA-206 overexpression. The expression of SOD1 was repressed by miRNA-206 overexpression, indicating SOD1 as a direct target of miRNA-206. Overexpression of miRNA-206 increased reactive oxygen species (ROS) levels *in vitro* and *in vivo*. Thus, overexpression of miRNA-206 promoted ANR by targeting SOD1 and production of ROS.

It was also reported that miRNA-206 overexpression in an experimental canine model inhibited expression of the *GCH1* gene, which encodes GTP cyclohydrolase I, a key enzyme in *de novo* synthesis of tetrahydrobiopterin (BH4), an essential cofactor for NO synthesis.⁽⁷⁸⁾ Previous studies reported that increased tetrahydrobiopterin (BH4) and NO content negatively regulated nerve regeneration. These findings suggested that *GCH1* downregulation exacerbates ANR by decreasing atrial BH4 and NO content modulated by miRNA-206 in A-TP canines.

Dysregulation of Ca^{2+} level in AF mediated by miRNAs

Calcium plays a fundamental role in the pathophysiology of AF. Enhanced SR Ca^{2+} leak through ryanodine receptor type-2 (RyR2), in combination with larger I(NCX) for a given (SR) Ca^{2+} release and increased diastolic [Ca^{2+}] (i)-voltage coupling gain, causes AF-promoting atrial delayed afterdepolarizations/triggered activity in AF patients.⁽⁷⁹⁾ The rise in intracellular calcium concentrations also activates pro-fibrotic pathways leading to SR in the atria that produces a substrate that can support the complex re-entry seen in AF.⁽⁸⁰⁾ Ca^{2+} -related miRNAs have been found to be significant pathophysiological contributors in conditions like myocardial ischemic injury, cardiac hypertrophy, heart failure, ventricular arrhythmogenesis, and AF.⁽⁸¹⁾

miRNA-106

Recent studies have shown that the level of RyR2 protein is elevated in atria of patients with paroxysmal AF, suggesting that miRNA-mediated post-transcriptional regulation of RyR2 might be an underlying mechanism. Bioinformatic analysis suggests that miRNA-106b, a member of the miRNA-106b-25 cluster, could bind to RyR2-3'-untranslated region and suppress its translation. Quantitative real-time PCR showed that the level of mature miRNA-106b was lower in atria of patients with paroxysmal AF than in patients in sinus rhythm. Ca^{2+} -spark frequency and total sarcoplasmic reticulum Ca^{2+} -leak were increased in atrial myocytes of miRNA-106b-25(-/-) mice, and these mice exhibited more frequent atrial ectopy and were also more susceptible to pacing-induced AF than wild-type littermates.⁽⁸²⁾

miRNA-208

Results from analysis of miRNA-208b-overexpressing HL-1 atrial myocytes and from myocytes isolated from CAF patients showed that aberrant miRNA-208b levels reduced the expression and function of L-type Ca²⁺ channel subunits (CACNA1C and CACNB2) as well as the sarcoplasmic reticulum-Ca²⁺ pump SERCA2.⁽⁸³⁾

The above data on the role of miRNAs in the pathophysiology of AF and the possibility of being used as AF biomarkers are summarized in Table 1.

miRNAs as AF biomarkers

miRNA expression in tissue and blood can be used as a biomarker for various diseases. A biomarker has been defined as “a characteristic that is objectively measured and evaluated as an indicator of normal biologic processes, pathogenic process, or pharmacologic responses to a therapeutic intervention”⁽⁸⁴⁾ and “A defined characteristic that is measured as an indicator of normal biological processes, pathogenic processes, or responses to an exposure or intervention, including therapeutic interventions.”⁽⁸⁵⁾

Currently, there are no suitable biomarkers for the primary diagnosis of AF. The use of natriuretic peptides and troponins, which are of very high value in myocardial infarction and HF, is ineffective with regard to the diagnosis and prognosis of AF.⁽⁶⁾ High stability, sensitivity, specificity, and prognostic properties of

c-miRNAs make them an attractive biomarker for the early diagnosis of numerous diseases, including CVD.⁽⁸⁶⁾ Moreover, miRNA expression can be detected both in biological fluids (plasma, serum), and in individual heart tissues and blood cells.

c-miRNAs exhibit numerous essential characteristics of biomarkers: While they are extremely stable in circulation, their expression is tissue-/disease-specific, and they can be easily detected using sequence-specific amplification methods. These features of c-miRNAs are helpful in the development of non-invasive assays to monitor the progress of CVDs.^(87,88)

Studies performed by Natsume et al.⁽⁸⁸⁾ revealed that four miRNAs (miRNA-99a-5p, miRNA-192-5p, miRNA-214-3p, and miRNA-342-5p) were significantly upregulated in AF patients. A receiver-operating characteristics curve indicated that miRNA-214-3p and miRNA-342-5p had the highest accuracy. The combination of the four miRNAs modestly improved the predictive accuracy for AF (76% sensitivity, 80% specificity).

c-miRNAs might be potential biomarkers of AF. They are easily accessible, relatively stable and in some instances tissue specific. At the least, c-miRNAs may offer new insights into the mechanisms of AF.⁽⁸⁹⁾ Further studies are required to gain a better understanding of the role of miRNAs in the pathophysiology of AF, which will provide tremendous opportunities for targeted therapy of AF in the near future.

Table 1.

The role of miRNAs in the pathophysiology of AF and the possibility of being used as AF biomarkers

Pathophysiological mechanism	miRNA	Targets and effects	A relationship of the elevated/decreased levels of miRNAs with AF	Source
Electrical remodeling	miRNA-1	Downregulation of the expression of the KCNE1, KCNB2, HCN2, and HCN4 genes	Both increased and decreased miRNA-1 expression may be associated with AF	[42-45]
	miRNA-328	Downregulation of the expression of the CACNA1C and CACNB1 genes	An increased miRNA-328 expression is associated with AF	[48-50]
	miRNA-499	Downregulation of the expression of the KCNN3 and CACNB2 genes	An increased miRNA-499 expression is associated with AF	[51-53]
Structural remodeling	miRNA-21	Downregulation of the expression of the CACNA1C and CACNB2 mRNAs	An increased miRNA-21 expression is associated with AF	[56-58]
	miRNA-29	Downregulation of the expression of the COL1A1, COL3A1, and FBN1 genes	Down-expression of miRNA-29 is associated with AF	[62, 63]
	miRNA-126	Downregulation of the EGFL7 expression	Down-expression of miRNA-126 is associated with AF	[64,65]
	miRNA-483	Upregulation of the expression of the IGF2 gene	An increased miRNA-483 expression is associated with AF	[70]
Remodeling of ANS	miRNA-30	Downregulation of the expression of IKACH gene	An increased miRNA-30 expression is associated with AF	[76]
	miRNA-206	Downregulation of the expression of the SOD1 и GCH1 genes	An increased miRNA-206 expression is associated with AF	[77,78]
Dysregulation of Ca ²⁺ level	miRNA-106	Downregulation of the expression of the RYR2 gene	Down-expression of miRNA-106 is associated with AF	[82]
	miRNA-208	Downregulation of the expression of the SERCA2a, CACNA1C, and CACNB2 genes	An increased miRNA-208 expression is associated with AF	[83]

Competing Interests

The authors declare that they have no competing interests.

References

1. Ko D, Rahman F, Martins MA, Hylek EM, Ellinor PT, Schnabel RB, Benjamin EJ, et al. Atrial fibrillation in women: treatment. *Nat Rev Cardiol*. 2017;14(2):113-124. doi: 10.1038/nrcardio.2016.171.
2. Pellman J, Sheikh F. Atrial fibrillation: mechanisms, therapeutics, and future directions. *Compr Physiol*. 2015;5(2):649-65. doi: 10.1002/cphy.c140047.
3. Turagam MK, Mirza M, Werner PH, Sra J, Kress DC, Tajik AJ, Jahangir A. Circulating Biomarkers Predictive of Postoperative Atrial Fibrillation. *Cardiol Rev*. 2016;24(2):76-87. doi: 10.1097/CRD.000000000000059.
4. Chaulin AM, Karslyan LS, Bazyuk EV, Nurbaltaeva DA, Duplyakov DV. [Clinical and Diagnostic Value of Cardiac Markers in Human Biological Fluids]. *Kardiologiya*. 2019;59(11):66-75. doi: 10.18087/cardio.2019.11.n414
5. Chaulin AM, Duplyakov DV. Increased cardiac troponins, not associated with acute coronary syndrome. Part 1. *Kardiologiya: novosti, mneniya, obuchenie* [Cardiology: News, Opinions, Training]. 2019;7(2):13-23. doi: 10.24411/2309-1908-2019-12002.
6. Chaulin AM, Duplyakov DV. Increased cardiac troponins, not associated with acute coronary syndrome. Part 2. *Kardiologiya: novosti, mneniya, obuchenie* [Cardiology: News, Opinions, Training]. 2019;7(2):24-35. doi: 10.24411/2309-1908-2019-12003.
7. Mahajan N, Mehta Y, Rose M, Shani J, Lichstein E. Elevated troponin level is not synonymous with myocardial infarction. *Int J Cardiol*. 2006;111(3):442-9. doi:10.1016/j.ijcard.2005.08.029
8. Gonzalez-Del-Hoyo M, Cediell G, Carrasquer A, Bonet G, Vasquez-Nunez K, Boque C, et al. Prognostic implications of troponin I elevation in emergency department patients with tachyarrhythmia. *Clin Cardiol*. 2019;42(5):546-552. doi: 10.1002/clc.23175.
9. Zellweger MJ, Schaer BA, Cron TA, Pfisterer ME, Osswald S. Elevated troponin levels in the absence of coronary artery disease after supraventricular tachycardia. *Swiss Med Wkly*. 2003;133(31-32):439-441. doi: 2003/31/smw-10288
10. Lee SH, Park SJ, Byeon K, On YK, Kim JS, Shin DG, et al. Risk factors between patients with lone and non-lone atrial fibrillation [published correction appears in *J Korean Med Sci*. 2014 Aug;29(8):1183]. *J Korean Med Sci*. 2013;28(8):1174-1180. doi:10.3346/jkms.2013.28.8.1174
11. Fox CS, Parise H, D'Agostino RB Sr, Lloyd-Jones DM, Vasan RS, Wang TJ, et al. Parental atrial fibrillation as a risk factor for atrial fibrillation in offspring. *JAMA*. 2004;291(23):2851-2855. doi: 10.1001/jama.291.23.2851.
12. Low SK, Takahashi A, Ebana Y, Ozaki K, Christophersen IE, Ellinor PT, et al. Identification of six new genetic loci associated with atrial fibrillation in the Japanese population. *Nat Genet*. 2017;49(6):953-958. doi: 10.1038/ng.3842.
13. Lee RC, Feinbaum RL, Ambros V. The *C. elegans* heterochronic gene *lin-4* encodes small RNAs with antisense complementarity to *lin-14*. *Cell*. 1993;75(5):843-854. doi: 10.1016/0092-8674(93)90529-y.
14. Tufekci KU, Meuwissen RL, Genc S. The role of microRNAs in biological processes. *Methods Mol Biol*. 2014;1107:15-31. doi:10.1007/978-1-62703-748-8_2
15. Ardekani AM, Naeini MM. The Role of MicroRNAs in Human Diseases. *Avicenna J Med Biotechnol*. 2010;2(4):161-79.
16. Schetter AJ, Heegaard NH, Harris CC. Inflammation and cancer: interweaving microRNA, free radical, cytokine and p53 pathways. *Carcinogenesis*. 2010;31(1):37-49. doi:10.1093/carcin/bgp272
17. Yu Z, Willmarth NE, Zhou J, Katiyar S, Wang M, Liu Y, et al. microRNA 17/20 inhibits cellular invasion and tumor metastasis in breast cancer by heterotypic signaling. *Proc Natl Acad Sci U S A*. 2010;107(18):8231-8236. doi:10.1073/pnas.1002080107
18. Liu C, Kelnar K, Liu B, Chen X, Calhoun-Davis T, Li H, et al. The microRNA miR-34a inhibits prostate cancer stem cells and metastasis by directly repressing CD44. *Nat Med*. 2011;17(2):211-215. doi:10.1038/nm.2284
19. Small EM, Olson EN. Pervasive roles of microRNAs in cardiovascular biology. *Nature*. 2011;469(7330):336-342. doi:10.1038/nature09783
20. Wang GK, Zhu JQ, Zhang JT, Li Q, Li Y, He J, et al. Circulating microRNA: a novel potential biomarker for early diagnosis of acute myocardial infarction in humans. *Eur Heart J*. 2010;31(6):659-666. doi:10.1093/eurheartj/ehq013
21. Contu R, Latronico MV, Condorelli G. Circulating microRNAs as potential biomarkers of coronary artery disease: a promise to be fulfilled? *Circ Res*. 2010;107(5):573-574. doi:10.1161/CIRCRESAHA.110.227983
22. Miller BH, Wahlestedt C. MicroRNA dysregulation in psychiatric disease. *Brain Res*. 2010;1338:89-99. doi:10.1016/j.brainres.2010.03.035
23. Haramati S, Chapnik E, Sztainberg Y, Eilam R, Zwang R, Gershoni N, et al. miRNA malfunction causes spinal motor neuron disease. *Proc Natl Acad Sci U S A*. 2010;107(29):13111-13116. doi:10.1073/pnas.1006151107
24. Leeper NJ, Cooke JP. MicroRNA and mechanisms of impaired angiogenesis in diabetes mellitus. *Circulation*. 2011;123(3):236-238. doi:10.1161/CIRCULATIONAHA.110.003855
25. Pandey AK, Agarwal P, Kaur K, Datta M. MicroRNAs in diabetes: tiny players in big disease. *Cell Physiol Biochem*. 2009;23(4-6):221-232. doi:10.1159/000218169
26. Fulci V, Scappucci G, Sebastiani GD, Giannitti C, Franceschini D, Meloni F, et al. miR-223 is overexpressed in T-lymphocytes of patients affected by rheumatoid arthritis. *Hum Immunol*. 2010;71(2):206-211. doi:10.1016/j.humimm.2009.11.008
27. Mattes J, Collison A, Plank M, Phipps S, Foster PS. Antagonism of microRNA-126 suppresses the effector function of TH2 cells and the development of allergic airways disease. *Proc Natl Acad Sci U S A*. 2009;106(44):18704-18709. doi:10.1073/pnas.0905063106
28. Polikepahad S, Knight JM, Naghavi AO, Opl T, Creighton CJ, Shaw C, et al. Proinflammatory role for let-7 microRNAs in experimental asthma. *J Biol Chem*. 2010;285(39):30139-30149. doi:10.1074/jbc.M110.145698
29. van Rooij E, Sutherland LB, Liu N, Williams AH, McAnally J, et al. A signature pattern of stress-responsive microRNAs that can evoke cardiac hypertrophy and heart failure. *Proc Natl Acad Sci U S A*. 2006;103(48):18255-18260. doi:10.1073/pnas.0608791103
30. Luo X, Yang B, Nattel S. MicroRNAs and atrial fibrillation: mechanisms and translational potential. *Nat Rev Cardiol*. 2015;12(2):80-90. doi:10.1038/nrcardio.2014.178
31. Arora P, Wu C, Khan AM, Bloch DB, Davis-Dusenbery BN, Ghorbani A, et al. Atrial natriuretic peptide is negatively

- regulated by microRNA-425. *J Clin Invest.* 2013;123(8):3378-3382. doi:10.1172/JCI67383
32. van Rooij E, Sutherland LB, Thatcher JE, DiMaio JM, Naseem RH, Marshall WS, et al. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc Natl Acad Sci U S A.* 2008;105(35):13027-13032. doi:10.1073/pnas.0805038105
33. Girmatsion Z, Biliczki P, Bonauer A, Wimmer-Greinecker G, Scherer M, Moritz A, et al. Changes in microRNA-1 expression and IK1 up-regulation in human atrial fibrillation. *Heart Rhythm.* 2009;6(12):1802-1809. doi:10.1016/j.hrthm.2009.08.035
34. Cardin S, Guasch E, Luo X, Naud P, Quang KL, Shi YF, et al. Role for microRNA-21 in atrial profibrillatory fibrotic remodeling associated with experimental postinfarction heart failure. *Circ Arrhythm Electrophysiol.* 2012;5(5):1027-1035. doi:10.1161/CIRCEP.112.973214
35. Duisters RF, Tijssen AJ, Schroen B, Leenders JJ, Lentink V, van der Made I, et al. miR-133 and miR-30 Regulate Connective Tissue Growth Factor. *Circ Res.* 2009;104(2):170-178. doi:10.1161/CIRCRESAHA.108.182535
36. Hodgkinson CP, Kang MH, Dal-Pra S, Mirosos M, Dzau VJ. MicroRNAs and Cardiac Regeneration. *Circ Res.* 2015;116(10):1700-1711. doi:10.1161/CIRCRESAHA.116.304377
37. van Rooij E, Olson EN. MicroRNA therapeutics for cardiovascular disease: opportunities and obstacles. *Nat Rev Drug Discov.* 2012;11(11):860-872. doi:10.1038/nrd3864
38. Weckbach LT, Grabmaier U, Clauss S, Wakili R. MicroRNAs as a diagnostic tool for heart failure and atrial fibrillation. *Curr Opin Pharmacol.* 2016;27:24-30. doi:10.1016/j.coph.2016.01.001
39. van den Berg NWE, Kawasaki M, Berger WR, Neefs J, Meulendijks E, Tijssen AJ, de Groot JR. MicroRNAs in Atrial Fibrillation: from Expression Signatures to Functional Implications. *Cardiovasc Drugs Ther.* 2017;31(3):345-365. doi:10.1007/s10557-017-6736-z
40. Chaldoupi SM, Loh P, Hauer RN, de Bakker JM, van Rijen HV. The role of connexin40 in atrial fibrillation. *Cardiovasc Res.* 2009;84(1):15-23. doi:10.1093/cvr/cvp203
41. Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, et al. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet.* 2006;38(2):228-233. doi:10.1038/ng1725
42. Jia X, Zheng S, Xie X, Zhang Y, Wang W, Wang Z, et al. MicroRNA-1 accelerates the shortening of atrial effective refractory period by regulating KCNE1 and KCNB2 expression: an atrial tachypacing rabbit model. *PLoS One.* 2013;8(12):e85639. doi:10.1371/journal.pone.0085639
43. Tsoporis JN, Fazio A, Rizos IK, Izhar S, Proteau G, Salpeas V, et al. Increased right atrial appendage apoptosis is associated with differential regulation of candidate MicroRNAs 1 and 133A in patients who developed atrial fibrillation after cardiac surgery. *J Mol Cell Cardiol.* 2018;121:25-32. doi:10.1016/j.yjmcc.2018.06.005
44. Li YD, Hong YF, Yusufuaji Y, Tang BP, Zhou XH, Xu GJ, et al. Altered expression of hyperpolarization-activated cyclic nucleotide-gated channels and microRNA-1 and -133 in patients with age-associated atrial fibrillation. *Mol Med Rep.* 2015;12(3):3243-3248. doi:10.3892/mmr.2015.3831
45. Girmatsion Z, Biliczki P, Bonauer A, Wimmer-Greinecker G, Scherer M, Moritz A, et al. Changes in microRNA-1 expression and IK1 up-regulation in human atrial fibrillation. *Heart Rhythm.* 2009;6(12):1802-1809. doi:10.1016/j.hrthm.2009.08.035
46. Lu Y, Hou S, Huang D, Luo X, Zhang J, Chen J, Xu W. Expression profile analysis of circulating microRNAs and their effects on ion channels in Chinese atrial fibrillation patients. *Int J Clin Exp Med.* 2015;8(1):845-853. Published 2015 Jan 15.
47. Santulli G, Iaccarino G, De Luca N, Trimarco B, Condorelli G. Atrial fibrillation and microRNAs. *Front Physiol* 2014;5:15. doi:10.3389/fphys.2014.00015
48. Lu Y, Zhang Y, Wang N, Pan Z, Gao X, Zhang F, et al. MicroRNA-328 contributes to adverse electrical remodeling in atrial fibrillation. *Circulation.* 2010;122(23):2378-2387. doi:10.1161/CIRCULATIONAHA.110.958967
49. Kim GH. MicroRNA regulation of cardiac conduction and arrhythmias. *Transl Res.* 2013;161(5):381-392. doi:10.1016/j.trsl.2012.12.004
50. Soeki T, Matsuura T, Bando S, Tobiume T, Uematsu E, Ise T, et al. Relationship between local production of microRNA-328 and atrial substrate remodeling in atrial fibrillation. *J Cardiol.* 2016;68(6):472-477. doi:10.1016/j.jjcc.2015.12.007
51. Ling TY, Wang XL, Chai Q, Lau TW, Koestler CM, Park SJ, et al. Regulation of the SK3 channel by microRNA-499--potential role in atrial fibrillation. *Heart Rhythm.* 2013;10(7):1001-1009. doi:10.1016/j.hrthm.2013.03.005
52. Ling TY, Wang XL, Chai Q, Lu T, Stulak JM, Joyce LD, et al. Regulation of cardiac CACNB2 by microRNA-499: Potential role in atrial fibrillation. *BBA Clin.* 2017;7:78-84. doi:10.1016/j.bbacli.2017.02.002
53. da Silva AMG, de Araujo JNG, de Oliveira KM, Novaes AEM, Lopes MB, de Sousa JCV, et al. Circulating miRNAs in acute new-onset atrial fibrillation and their target mRNA network. *J Cardiovasc Electrophysiol.* 2018;29(8):1159-1166.
54. Xu J, Cui G, Esmailian F, Plunkett M, Marelli D, Ardehali A, et al. Atrial extracellular matrix remodeling and the maintenance of atrial fibrillation. *Circulation.* 2004;109(3):363-368. doi:10.1161/01.CIR.0000109495.02213.52
55. Liu H, Chen GX, Liang MY, Qin H, Rong J, Yao JP, Wu ZK. Atrial fibrillation alters the microRNA expression profiles of the left atria of patients with mitral stenosis. *BMC Cardiovasc Disord.* 2014;14:10. doi:10.1186/1471-2261-14-10
56. Barana A, Matamoros M, Dolz-Gaiton P, Pérez-Hernández M, Amoros I, Nunez M, et al. Chronic atrial fibrillation increases microRNA-21 in human atrial myocytes decreasing L-type calcium current. *Circ Arrhythm Electrophysiol.* 2014;7(5):861-868. doi:10.1161/CIRCEP.114.001709
57. Adam O, Lohfelme B, Thum T, Gupta SK, Puhl SL, Schafers HJ, et al. Role of miR-21 in the pathogenesis of atrial fibrosis. *Basic Res Cardiol.* 2012;107(5):278. doi:10.1007/s00395-012-0278-0
58. Thum T, Gross C, Fiedler J, Fischer T, Kissler S, Bussen M, et al. MicroRNA-21 contributes to myocardial disease by stimulating MAP kinase signalling in fibroblasts. *Nature.* 2008;456(7224):980-984. doi:10.1038/nature07511
59. Cantley LC. The phosphoinositide 3-kinase pathway. *Science.* 2002;296(5573):1655-1657. doi:10.1126/science.296.5573.1655
60. Zhang K, Zhao L, Ma Z, Wang W, Li X, Zhang Y, et al. Doxycycline Attenuates Atrial Remodeling by Interfering with MicroRNA-21 and Downstream Phosphatase and Tensin Homolog (PTEN)/Phosphoinositide 3-Kinase (PI3K) Signaling Pathway. *Med Sci Monit.* 2018;24:5580-5587. doi:10.12659/MSM.909800
61. van Rooij E, Sutherland LB, Thatcher JE, DiMaio JM, Naseem RH, Marshall WS, et al. Dysregulation of microRNAs after myocardial infarction reveals a role of miR-29 in cardiac fibrosis. *Proc Natl Acad Sci U S A.* 2008;105(35):13027-13032.

62. Kriegel AJ, Liu Y, Fang Y, Ding X, Liang M. The miR-29 family: genomics, cell biology, and relevance to renal and cardiovascular injury. *Physiol Genomics*. 2012;44(4):237-244. doi: 10.1152/physiolgenomics.00141.2011.
63. Dawson K, Wakili R, Ordog B, Clauss S, Chen Y, Iwasaki Y, et al. MicroRNA29: a mechanistic contributor and potential biomarker in atrial fibrillation. *Circulation*. 2013;127(14):1466-1475, 1475e1-28. doi: 10.1161/CIRCULATIONAHA.112.001207.
64. Liakouli V, Cipriani P, Di Benedetto P, Panzera N, Ruscitti P, Pantano I, et al. Epidermal Growth Factor Like-domain 7 and miR-126 are abnormally expressed in diffuse Systemic Sclerosis fibroblasts. *Sci Rep*. 2019;9(1):4589. Published 2019 Mar 14. doi:10.1038/s41598-019-39485-8
65. Wei XJ, Han M, Yang FY, Wei GC, Liang ZG, Yao H, et al. Biological significance of miR-126 expression in atrial fibrillation and heart failure. *Braz J Med Biol Res*. 2015;48(11):983-989. doi: 10.1590/1414-431X20154590.
66. McManus DD, Tanriverdi K, Lin H, Esa N, Kinno M, Mandapati D, et al. Plasma microRNAs are associated with atrial fibrillation and change after catheter ablation (the miRhythm study). *Heart Rhythm*. 2015;12(1):3-10. doi: 10.1016/j.hrthm.2014.09.050.
67. Goren Y, Meiri E, Hogan C, Mitchell H, Lebanony D, Salman N, et al. Relation of reduced expression of MiR-150 in platelets to atrial fibrillation in patients with chronic systolic heart failure. *Am J Cardiol*. 2014;113(6):976-981. doi: 10.1016/j.amjcard.2013.11.060.
68. Lutter D, Marr C, Krumsiek J, Lang EW, Theis FJ. Intronic microRNAs support their host genes by mediating synergistic and antagonistic regulatory effects. *BMC Genomics*. 2010;11:224. Published 2010 Apr 6. doi:10.1186/1471-2164-11-224
69. Qiao Y, Ma N, Wang X, Hui Y, Li F, Xiang Y et al. MiR-483-5p controls angiogenesis in vitro and targets serum response factor. *FEBS Lett*. 2011;585(19):3095-3100. doi:10.1016/j.febslet.2011.08.039
70. Harling L, Lambert J, Ashrafian H, Darzi A, Gooderham NJ, Athanasiou T. Elevated serum microRNA 483-5p levels may predict patients at risk of post-operative atrial fibrillation. *Eur J Cardiothorac Surg*. 2017;51(1):73-78. doi: 10.1093/ejcts/ezw245.
71. Wang YC, Hu YW, Sha YH, Gao JJ, Ma X, Li SF, et al. Ox-LDL Upregulates IL-6 Expression by Enhancing NF- κ B in an IGF2-Dependent Manner in THP-1 Macrophages. *Inflammation*. 2015;38(6):2116-2123. doi:10.1007/s10753-015-0194-1
72. Chen PS, Chen LS, Fishbein MC, Lin SF, Nattel S. Role of the autonomic nervous system in atrial fibrillation: pathophysiology and therapy. *Circ Res*. 2014;114(9):1500-1515. doi:10.1161/CIRCRESAHA.114.303772
73. Patterson E, Po SS, Scherlag BJ, Lazzara R. Triggered firing in pulmonary veins initiated by in vitro autonomic nerve stimulation. *Heart Rhythm*. 2005;2(6):624-631. doi:10.1016/j.hrthm.2005.02.012
74. Rao M, Hu J, Zhang Y, Gao F, Zhang F, Yang Z, et al. Time-dependent cervical vagus nerve stimulation and frequency-dependent right atrial pacing mediates induction of atrial fibrillation. *Anatol J Cardiol*. 2018;20(4):206-212. doi: 10.14744/AnatolJCardiol.2018.73558.
75. Shen MJ, Choi EK, Tan AY, Lin SF, Fishbein MC, Chen LS, Chen PS. Neural mechanisms of atrial arrhythmias. *Nat Rev Cardiol*. 2011;9(1):30-39. doi: 10.1038/nrcardio.2011.139.
76. Morishima M, Iwata E, Nakada C, Tsukamoto Y, Takanari H, Miyamoto S, et al. Atrial Fibrillation-Mediated Upregulation of miR-30d Regulates Myocardial Electrical Remodeling of the G-Protein-Gated K(+) Channel, IK₁ACH. *Circ J*. 2016;80(6):1346-1355. doi: 10.1253/circj.CJ-15-1276.
77. Zhang Y, Zheng S, Geng Y, Xue J, Wang Z, Xie X, et al. MicroRNA profiling of atrial fibrillation in canines: miR-206 modulates intrinsic cardiac autonomic nerve remodeling by regulating SOD1. *PLoS One*. 2015;10(3):e0122674. doi: 10.1371/journal.pone.0122674.
78. Wei J, Zhang Y, Li Z, Wang X, Chen L, Du J, et al. GCH1 attenuates cardiac autonomic nervous remodeling in canines with atrial-tachypacing via tetrahydrobiopterin pathway regulated by microRNA-206. *Pacing Clin Electrophysiol*. 2018;41(5):459-471. doi: 10.1111/pace.13289.
79. Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, et al. Enhanced sarcoplasmic reticulum Ca²⁺ leak and increased Na⁺-Ca²⁺ exchanger function underlie delayed afterdepolarizations in patients with chronic atrial fibrillation. *Circulation*. 2012;125(17):2059-2070. doi:10.1161/CIRCULATIONAHA.111.067306
80. Denham NC, Pearman CM, Caldwell JL, Madders GWP, Eisner DA, Trafford AW, Dibb KM. Calcium in the Pathophysiology of Atrial Fibrillation and Heart Failure. *Front Physiol*. 2018;9:1380. doi: 10.3389/fphys.2018.01380.
81. Harada M, Luo X, Murohara T, Yang B, Dobrev D, Nattel S. MicroRNA regulation and cardiac calcium signaling: role in cardiac disease and therapeutic potential. *Circ Res*. 2014;114(4):689-705. doi:10.1161/CIRCRESAHA.114.301798.
82. Chiang DY, Kongchan N, Beavers DL, Alsina KM, Voigt N, Neilson JR, et al. Loss of microRNA-106b-25 cluster promotes atrial fibrillation by enhancing ryanodine receptor type-2 expression and calcium release. *Circ Arrhythm Electrophysiol*. 2014;7(6):1214-1222. doi:10.1161/CIRCEP.114.001973
83. Canon S, Caballero R, Herraiz-Martínez A, Perez-Hernández M, Lopez B, Atienza F, et al. miR-208b upregulation interferes with calcium handling in HL-1 atrial myocytes: Implications in human chronic atrial fibrillation. *J Mol Cell Cardiol*. 2016;99:162-173. doi: 10.1016/j.yjmcc.2016.08.012.
84. Downing GL, NIH Definitions Working Group. Biomarkers and surrogate endpoints in clinical research: definitions and conceptual model. In *Biomarkers and Surrogate Endpoints: Clinical Research and Applications*, 1–9. Elsevier, Amsterdam, UK; 2000.
85. BEST (Biomarkers, EndpointS, and Other Tools) Resource [Internet]. FDA-NIH Biomarker Working Group. Silver Spring (MD): Food and Drug Administration (US); Bethesda (MD): National Institutes of Health (US); 2016. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK338448/>
86. D'Alessandra Y, Devanna P, Limana F, Straino S, Di Carlo A, Brambilla PG, et al. Circulating microRNAs are new and sensitive biomarkers of myocardial infarction. *Eur Heart J*. 2010;31:2765-2773. doi: 10.1093/eurheartj/ehq167
87. Felekakis K, Papanephytous C. Challenges in Using Circulating Micro-RNAs as Biomarkers for Cardiovascular Diseases. *Int J Mol Sci*. 2020;21(2):561. Published 2020 Jan 15. doi:10.3390/ijms21020561
88. Natsume Y, Oaku K, Takahashi K, Nakamura W, Oono A, Hamada S, et al. Combined Analysis of Human and Experimental Murine Samples Identified Novel Circulating MicroRNAs as Biomarkers for Atrial Fibrillation. *Circ J*. 2018;82(4):965-973. doi: 10.1253/circj.CJ-17-1194.
89. Zampetaki A, Willeit P, Drozdov I, Kiechl S, Mayr M. Profiling of circulating microRNAs: from single biomarkers to re-wired networks. *Cardiovasc Res*. 2012; 93 (4): 555-562. doi: 10.1093/cvr/cvr266.