

# Association between Hypodontia of the Permanent Lateral Incisors and other Dental Anomalies in School Children Aged 12-16 Years in Kosovo

Albena Reshitaj<sup>1\*</sup>; Krenar Reshitaj<sup>2</sup>; Erik Musliu<sup>3</sup>; Kanita Hoxha<sup>3</sup>

<sup>1</sup>Department of Orthodontics, Medical Faculty School of Dentistry, University of Prishtina

<sup>2</sup>Medical Faculty, University Clinical Center of Kosovo

<sup>3</sup>Department of Dentistry, Rezonanca Medical College  
Prishtina, Kosovo

## Abstract

**Background:** Hypodontia of lateral incisors (LI) is frequently associated with other dental anomalies. The objective of this study was to determine the association of LI with other dental anomalies by comparing the two groups: Group 1 with hypodontia of the maxillary LI (MLI1) and Group 2 with hypodontia of the mandibular LI (MLI2), in secondary school education students in Kosovo.

**Methods and Results:** A total of 3306 secondary school students aged 12-16 years, regardless of gender, were included in this prospective study. The abnormalities investigated were recorded by RTG-panoramic and dental charts. The teeth were recorded as a congenital absence when the mineralization of the crown, identified by panoramic tomography, was absent. In a sample of 3306 subjects, 77(2.3%) subjects were diagnosed with hypodontia. The highest percentage of hypodontia was found in the upper left LI in 20.4% of cases, in the upper right LI in 18.4% of cases, while the percentage of hypodontia of the lower left LI was 0.7% and 2.0% on the right side. A lower percentage of 0.7% or just one missing tooth was found in teeth 13, 32, and 46. The prevalence of LI hypodontia was as follows: MLI1 (Group 1) included 36 cases (92.3%) and MLI2 (Group 2) included only 3 cases (7.6%), which indicates a much higher percentage of cases with hypodontia of MLI1. In Group 1, 21(58.3%) cases of LI hypodontia were bilateral and 15(41.7%) unilateral; in Group 2, 2(66.7%) cases were unilateral and 1(33.3%) case – bilateral.

Among dental anomalies, the occurrence of rotation was found in 19(48.7%) cases with LI hypodontia: 47.2% cases in Group 1 and 66.7% cases in Group 2. The prevalence of dental inclination anomaly was 30.77% of all cases with hypodontia of LI: 27.8% of cases in Group 1 and 66.7% of cases in Group 2. The prevalence of ectopy was in 17.9% of cases of all hypodontia cases of LI: 16.7% of cases in Group 1 and 33.3% of cases in Group 2. Crown anomalies were evident in 7(17.9%) patients of all hypodontia cases of LI, all of which were in Group 1. Other anomalies such as microdontia were evident in two patients in Group 1. Transposition, bodily movement, and superposition were present in one patient in Group 1; in Group 2, one patient had transposition. At the same time, the frequency of dental transposition was significantly higher in Group 2 than in Group 1 ( $P=0.0209$ ). Other dental anomalies of crown and root and infraposition were not present in both groups. There were no significant differences in the other dental anomalies between the two groups.

**Conclusion:** The consequences of hypodontia in dental arches are obvious. Knowing the prevalence of hypodontia and its association with other dental anomalies helps classify the need for further treatment for the patients, whether orthodontic, prosthetic, or surgical. (**International Journal of Biomedicine. 2022;12(2):273-278.**)

**Key Words:** hypodontia • lateral incisors • dental anomalies

**For citation:** Reshitaj A, Reshitaj K, Musliu E, Hoxha K. Association between Hypodontia of the Permanent Lateral Incisors and other Dental Anomalies in School Children Aged 12-16 Years in Kosovo. International Journal of Biomedicine. 2022;12(2):273-278. doi:10.21103/Article12(2)\_OA13

## Introduction

Hypodontia is one of the most common developmental anomalies in humans,<sup>(1,2)</sup> which is characterized by the absence of one or more deciduous or permanent teeth, as well as on both upper and lower dentitions.<sup>(3)</sup> Hypodontia can cause masticatory dysfunction, impact speech, create aesthetic problems with

psychological implications affecting self-esteem,<sup>(4)</sup> and cause the need for orthodontic treatment and prosthetic compensation in some cases. Hypodontia of lateral incisors (LI) is frequently associated with other dental anomalies.

A variety of terminology describes this anomaly in the literature, such as tooth aplasia, congenitally missing teeth, missing teeth, dental agenesis, oligodontia, or anodontia.

The appearance of dental agenesis in permanent dentition varies. Polder et al., using meta-analysis, found that the prevalence of missing permanent teeth varies from 2.2% to 10.1%, excluding third molars, which are missing in about 20% of the general population.<sup>(5)</sup> This anomaly has multifactorial etiology, including genetic factors, phylogenetic and environmental factors,<sup>(6,7)</sup> and has variability in the size of the teeth, most often characterized by microdontia.<sup>(8)</sup>

The absence of anterior teeth and disorders regarding the position of the other teeth cause malocclusions that directly affect both function and aesthetics.<sup>(9-11)</sup>

The appearance of unilateral hypodontia in LI is, in most cases, accompanied by changes in the shape and size (peg-shaped) of the same tooth on the opposite side of the jaw. This means that both abnormalities deviate from the same genes, but the difference lies in changes in gene expressiveness. In people with permanent dental hypodontia, the persistence of a deciduous tooth is often seen as a result of the absence of a permanent tooth. This includes the second deciduous mandibular molars and deciduous maxillary canines. In the absence of a permanent upper lateral incisor, the canine migrates forward and emerges between the central incisor and the deciduous canine.<sup>(12)</sup>

Hypodontia in deciduous teeth is a relatively rare phenomenon. The prevalence is from 0.1% to 0.9%.<sup>(13)</sup> This anomaly is more often localized in the maxilla than in the mandible, while the missing teeth are usually the upper LI. Hypodontia of a deciduous tooth in primary dentition can also be associated with hypodontia of the same tooth in permanent dentition.<sup>(14)</sup>

The persistence of deciduous teeth often occurs in association with hypodontia. As a result, when a tooth is missing, it is thought that there is a lack of impulse for resorption of the root of the deciduous tooth, which results in the tooth remaining in the jaw for a long time. We often notice the persistence of deciduous canines in cases with hypodontia. As a result of hypodontia, deciduous teeth are prevalent because their root resorption is not helped due to the missing permanent teeth. According to some authors, the retention of primary teeth prevents the vertical development of the alveolar process.<sup>(15,16)</sup>

The prevalence of hypodontia of permanent maxillary LI ranges from 6% to 8% in different ethnic groups, and molecular genetics has identified shared genetic mutations in families with tooth agenesis.<sup>(17)</sup> Hypodontia of permanent maxillary LI is also associated with other dental anomalies, while the same genetic mutation may have a variable phenotypical expression.<sup>(18)</sup>

There have also been suggestions of various inheritance models related to this anomaly, including dominant autosomal incomplete penetration, recessive autosomal, gender-linked inheritance, and polygenic inheritance model. These genes are associated with about 120 syndromes, such as cleft lip, cleft palate, and ectodermal dysplasia, as well as Down, Rieger, and Book Syndromes.<sup>(19)</sup>

Hypodontia of LI is not an isolated phenomenon because they are often reported associated with other dental anomalies, such as LI in the form of a peg, transposition, delayed tooth

development, ectopic eruption, stagnant deciduous teeth, inclination, infraposition, and other anomalies in the size and shape of the teeth.<sup>(12)</sup> A study of orthodontic patients with at least one canine in the palatal ectopic position showed that LI near these canines were missing in a high percentage of cases.<sup>(20)</sup>

Peg-shaped maxillary LI were found in 5.5% of family members (proband) with hypodontia, compared with a frequency of 1.7% of the general population.<sup>(21)</sup> In a study by Alvesalo and Portin, the frequency and inheritance models were researched, and it was observed that the peg-shaped upper LI and the mesiodistal reduction of the upper LI are strongly different expressions of the autosomal dominant gene with reduced penetration.<sup>(22)</sup>

Hypodontia has a strong connection with microdontia. Congenital absence of LI is often associated with a reduced tooth on the contralateral side with genetic etiology that, combined with hypodontia, can result in multiple diastemas and rotations of adjacent teeth.<sup>(23)</sup> Previous studies have reported an association between hypodontia of maxillary incisors and other dental anomalies such as transposition,<sup>(24)</sup> displacement of palatal canines,<sup>(25)</sup> and premolar rotation.<sup>(26)</sup>

The studies about the differences between hypodontia of maxillary and mandibular LI regarding their association with other dental anomalies are scarce. In this study, we aimed to investigate whether hypodontia of the maxillary LI has a higher prevalence in the presence of other dental abnormalities compared to the hypodontia group of mandibular LI. This research may provide further evidence on the field-specific genetic control of tooth development.

The objective of this study was to determine the association of LI with other dental anomalies by comparing the two groups: Group 1 with hypodontia of the maxillary LI (MLI1) and Group 2 with hypodontia of the mandibular LI (MLI2), in secondary school education students in Kosovo.

## Materials and Methods

A total of 3306 secondary school students aged 12-16 years, regardless of gender, were included in this prospective study. All students attended their respective secondary schools throughout Kosovo. The selection of schools and participants was random, involving all regions of the country, both rural and urban. Examinations were performed by orthodontics and dentistry specialists. The abnormalities investigated were recorded by RTG-panoramic and dental charts. The teeth were recorded as a congenital absence when the mineralization of the crown, identified by panoramic tomography, was absent. The excluding criteria were a history of tooth loss from trauma, caries, periodontal disease, or orthodontic extractions.

The order of patients, based on age, was therefore selected by taking into account the delayed development of second mandibular premolars in boys<sup>(24)</sup> and according to the dental stage classification developed by Björk.<sup>(25)</sup>

Inclusion criteria were children of both genders from all regions and secondary schools throughout the country, subjects with all teeth present, students who have no previous history of trauma according to the anamnesis.

Exclusion criteria were students who provided data showing a previous history of tooth loss due to trauma, caries, periodontal disease, or orthodontic extractions; subjects with hypodontia associated with congenital syndromes or systemic diseases; all suspected cases in medical history and clinical examination.

The sample selection was made through the “cluster sampling” technique. This was based on the radiography verification of hypodontia, which was completed for all teeth, excluding the third molars from the study. The file for each subject was reviewed for medical histories, dental and family histories, and study models of maxillary and mandibular dental arches.

Statistical analysis was performed using statistical software package SPSS version 20.0 (Armonk, NY: IBM Corp.). Baseline characteristics were summarized as frequencies and percentages for categorical variables. The frequencies of categorical variables were compared using Pearson’s chi-squared test or Fisher’s exact test (2-tail), when appropriate. A two-proportion z-test was used to determine whether the two proportions were different from each other. A value of  $P < 0.05$  was considered significant.

This study was approved by the Ethics Committee of the Faculty of Medicine, the University of Prishtina, supported by the Ministry of Science and Education and the University Dental Clinical Center of Kosovo. Written informed consent was obtained from the parent/guardian/relative of each patient.

## Results

In a sample of 3306 subjects aged 12-16 years, 77(2.3%) subjects were diagnosed with hypodontia (Table 1).

**Table 1.**

**Presentation of cases frequency by gender, hypodontia and hypodontia in jaw, including LI hypodontia**

Gender	Number of cases, n (%)	Hypodontia frequency	Hypodontia in jaw			Hypodontia of LI	
			Maxillary	Mandibular	Both	Yes	No
Female	1566 (47.4%)	46 (2.9%)	26 (56.5%)	13 (28.3%)	7 (15.2%)	23 (1.47%)	23 (1.47%)
Male	1740 (52.6%)	31 (1.8%)	20 (64.5%)	8 (25.8%)	3 (9.7%)	16 (0.92%)	15 (0.86%)
Total	3306	77 (2.3%)	46 (59.7%)	21 (27.3%)	10 (13.0%)	39 (1.18%)	38 (1.15%)

Regarding the gender groups, 46(2.9%) were female and 31(1.80%) male. There were significant differences between the gender groups ( $z=2.2$ ,  $P=0.0278$ ). Among all 77 hypodontia cases, the presence of jaw-based hypodontia was 46(59.7%) cases in the maxillary jaw and 21(27.3%) cases in the mandibular jaw. Hypodontia of both jaws was present in 10(13.0%) cases. There were no significant differences between groups based on gender and jaw hypodontia. The prevalence of LI hypodontia was 1.18% or 39 cases in the total sample. The

presence of LI hypodontia was 39(50.6%) of overall hypodontia cases, of which 23(58.9%) were female and 16(41.3%) male.

The highest percentage of hypodontia was found in the upper left LI in 20.4% of cases, in the upper right LI in 18.4% of cases, while the percentage of hypodontia of the lower left LI was 0.7% and 2.0% on the right side (Table 2). A lower percentage of 0.7% or just one missing tooth was found in teeth 13, 32, and 46.

**Table 2.**

**Frequency of affected teeth (FDI notation) in hypodontia subjects**

Affected teeth	Number of cases	Percentage
12	27	18.4
13	1	0.7
14	10	6.8
15	6	4.1
22	30	20.4
23	4	2.7
24	11	7.5
25	5	3.4
31	3	2.0
32	1	0.7
34	7	4.8
35	12	8.2
41	4	2.7
42	3	2.0
44	10	6.8
45	12	8.2
46	1	0.7
Total	147	100

The prevalence of LI hypodontia was as follows: MLI1 (Group 1) included 36 cases (92.3%) and MLI2 (Group 2) included only 3 cases (7.6%), which indicates a much higher percentage of cases with hypodontia of MLI1 (Table 3). In Group 1, 21(58.3%) cases of hypodontia were bilateral and 15(41.7%) unilateral; in Group 2, 2(66.7%) cases were unilateral and 1(33.3%) case – bilateral.

Among dental anomalies, the occurrence of rotation was found in 19(48.7%) cases with LI hypodontia: 47.2% cases in Group 1 and 66.7% cases in Group 2. The prevalence of dental inclination anomaly was 30.77% of all cases with hypodontia of LI: 27.8% of cases in Group 1 and 66.7% of cases in Group 2. The prevalence of ectopy was in 17.9% of cases of all hypodontia cases of LI: 16.7% in Group 1 and 33.3% of cases in Group 2. Crown anomalies were evident in 7(17.9%) patients of all hypodontia cases of LI, all of which were in Group 1. Other anomalies such as microdontia were evident in two patients in Group 1. Transposition, bodily movement, and superposition were present in one patient in Group 1; in Group 2, one patient had transposition. Other dental anomalies of crown and root and infraposition were not present in both groups. At the same time, the frequency of dental transposition was significantly higher in Group 2 than in Group 1 ( $P=0.02088$ ). There were no significant differences in the other dental anomalies between the two groups.

**Table 3.**  
**The presentation of the frequency and distribution of dental anomalies in the group with hypodontia of MLI1 and MLI2**

Dental anomaly	Group 1 (Hypodontia of the MLI1)				Group 2 (Hypodontia of the MLI2)									
	Uni-lateral		Uni-lateral		Bila-teral		Total		Uni-lateral		Bila-teral		Total	
	n	%	n	%	n	%	n	%	n	%	n	%	n	%
	6	16.7	9	25.0	21	58.3	36	100	2	66.7	1	33.3	3	100
	12		22		12-22				42		32-42			
Rotation	4		6		7		17		2		0		2	
Inclination	3		3		4		10		2		0		2	
Bodily	0		0		1		1		0		0		0	
Superposition	0		1		0		1		0		0		0	
Infraposition	0		0		0		0		0		0		0	
Ectopia	2		4		0		6		1		0		1	
Transposition	0		1		0		1		1		0		1	
Crown anomaly	3		3		1		7		0		0		0	
Root anomaly	0		0		0		0		0		0		0	
Microdontia	1		0		1		2		0		0		0	

**Table 4.**  
**Distribution and prevalence of dental anomalies in groups MLI1 and MLI2**

Dental anomaly	Group 1 (MLI1) (n=36)	Group 2 (MLI2) (n=3)	Statistics	
	n (%)	n (%)	z-score	P-value
Rotation	17 (47.2)	2 (66.7)	-0.6474	0.5157
Inclination	10 (27.8)	2 (66.7)	-1.4022	0.16152
Bodily	1 (2.8)	0	0.2924	0.77182
Superposition	1 (2.8)	0	0.2924	0.77182
Infraposition	0	0	-	-
Ectopia	6 (16.7)	1 (33.3)	-0.7227	0.47152
Transposition	1 (2.8)	1 (33.3)	-2.3053	0.02088
Crown anomaly	7 (19.4)	0	0.8432	0.4009
Root anomaly	0	0	-	-
Microdontion	2 (5.6)	0	0.4191	0.67448

## Discussion

The data of our study, with a higher prevalence among females than males, follows the studies of Gokkaya & Kargul<sup>(27)</sup> and Badrov et al.<sup>(28)</sup> Also, the findings of the ratio of male to female, which was 1:1.4 in our study, is similar to the studies done by Baceti<sup>(26)</sup> and Muller et al.<sup>(29)</sup> but differ from the findings of Albashaireh & Khader,<sup>(30)</sup> Maklin et al.,<sup>(31)</sup> Rölling,<sup>(32)</sup> and Aasheim & Ogaard.<sup>(33)</sup>

The most common missing teeth are the LI of the upper jaw on the left side in 20.5% of cases and the right side in 17.8% of cases, followed by the second premolars of the lower

and upper jaw (10%), which is consistent with some studies.<sup>(11,34)</sup> The frequency of LI hypodontia in our study was higher in 50.6% of all cases with hypodontia, and similar results were found in the studies by Pinho et al.,<sup>(35)</sup> Celikoglu et al.,<sup>(36)</sup> and Silve Meza R.<sup>(37)</sup> The prevalence of hypodontia of maxillary LI was 46.7% of the total sample with hypodontia and is similar to the findings of a study by Al-Abdallah et al.<sup>(38)</sup>

In our study, the prevalence of uni- and bilateral maxillary LI hypodontia was 1.09% of the total sample and is consistent with other reported studies ranging from 0.3% to 11.3%.<sup>(3,37,39)</sup> According to a meta-analysis, the prevalence of agenesis of permanent teeth in the general population, excluding third molars, ranges from 3.2% to 7.6%.<sup>(34)</sup> This prevalence varies according to the tooth type. For example, hypodontia of maxillary LI was 1.7% of all samples, which is similar to our findings.

Bilateral agenesis of the maxillary LI occurs more frequently than unilateral agenesis.<sup>(40-43)</sup> Our study also shows a higher frequency of agenesis of bilateral maxillary LI in 56.4% of cases, but other studies by Pinho et al.<sup>(35)</sup> and Delli K et al.<sup>(44)</sup> showed less frequency of bilateral than unilateral agenesis.

A study by Celikoglu et al.<sup>(36)</sup> found significantly increased prevalence rates for ectopic eruption, transposition, and transmigration of the maxillary canines and reduced or peg-shaped maxillary LIs in their study sample. This is similar to the findings in our study, where the results indicate a high prevalence of dental anomalies, such as rotation, inclination, ectopic, and corona anomalies, in the group of patients with hypodontia of LI. Rotation of premolars was significantly associated with congenitally missing maxillary LI in the study by Baccetti.<sup>(26)</sup> This result was similar to the findings in our study, where we found a high prevalence of the dental rotation anomaly [19(48.7%) cases with LI hypodontia].

The lateral incisor of the maxilla was the tooth most often missing congenitally, as in the study of Augard & Gayard.<sup>(45)</sup> However, this is not consistent with the study of Al-Mulla et al.<sup>(46)</sup> who found that the second premolar of the mandibula is the most frequently missing tooth.

In a study about hypodontia, a critical issue is the patient's age at the time of the diagnosis, which tells us that the visibility of the dental germ on radiography depends on the stage of tooth mineralization.<sup>(43)</sup> The stages of tooth development are more closely related to tooth mineralization than the chronological age of tooth eruption.<sup>(47)</sup> Unilateral hypodontia is often associated with dysmorphia or microdontia corresponding to the contralateral tooth.<sup>(48)</sup> In another study by Pinho et al.,<sup>(49)</sup> the findings indicate that microdontia of the maxillary LI may represent a presentation of the molecular changes that lead to a developing defect of the maxillary lateral incisors. Therefore, considerable emphasis should be placed on the clinical diagnosis or family history where we may suspect missing teeth, and the treatment option for closing the spaces becomes unreal from the orthodontic aspect.<sup>(50)</sup>

Considering that the possibilities of treating patients with hypodontia are wide during treatment planning,<sup>(51)</sup> it is necessary to know the number of missing teeth, especially in the intercanine sector, the size of the teeth, the condition

of the teeth present, the condition of the periodontium, the position of the teeth, alveolar bone mass, general and local health status (soft tissue, lip line, gingival aesthetic condition, malocclusions, patient's age, the attitude of the child and parents towards the anomaly, occlusion, etiological factors, technical and financial possibilities of solving the problem).

Recognizing the prevalence and different models of hypodontia, strategies are created to treat this anomaly, which is not only statistically important but also important for the formation of an interdisciplinary professional team to cooperate in planning successful treatment.

The consequences of hypodontia in dental arches are obvious. Depending on the models of hypodontia, the need for the treatment ranges from simple to intensive. The problems that can occur in these patients are extensive, such that each case needs to be considered uniquely, in terms of approach and treatment. Knowing the prevalence of hypodontia and its association with other dental anomalies helps classify the need for further treatment for the patients, whether orthodontic, prosthetic, or surgical.

In our study, there were dental abnormalities associated with dental agenesis, which should be sought and investigated in patients with agenesis. It is particularly necessary, for those with agenesis of upper lateral incisors, and especially in young children for whom it is crucial that the orthodontist intercept these anomalies as early as possible, in order to establish proper treatment. The time-consuming and financial cost of extensive treatments of this anomaly is of interest to numerous clinical, basic science, and public health fields, such as orthodontics, pediatric dentistry, prosthodontics, periodontics, maxillofacial surgery, anatomy, anthropology, and even to the insurance companies.

## Competing Interests

The authors declare that they have no competing interests.

## References

1. Thind BS, Stirrups DR, Forgie AH, Larmour CJ, Mossey PA. Management of hypodontia: orthodontic considerations (II). *Quintessence Int.* 2005 May;36(5):345-53.
2. De Coster PJ, Marks LA, Martens LC, Huysseune A. Dental agenesis: genetic and clinical perspectives. *J Oral Pathol Med.* 2009 Jan;38(1):1-17. doi: 10.1111/j.1600-0714.2008.00699.x.
3. Fekonja A. Hypodontia in orthodontically treated children. *Eur J Orthod.* 2005 Oct;27(5):457-60. doi: 10.1093/ejo/cji027.
4. Oshagh M, Salehi P, Pakshir H, Bazayr L, Rakhshan V. Associations between normative and self-perceived orthodontic treatment needs in young-adult dental patients. *The Korean Journal of Orthodontics.* Korean Association of Orthodontists.2011;41(6):440-6. [Google Scholar]
5. Polder BJ, Van't Hof MA, Van der Linden FP, Kuijpers-Jagtman AM. A meta-analysis of the prevalence of dental agenesis of permanent teeth. *Community Dent Oral Epidemiol.* 2004 Jun;32(3):217-26. doi: 10.1111/j.1600-0528.2004.00158.x.
6. Vieira AR. Oral clefts and syndromic forms of tooth agenesis as models for genetics of isolated tooth agenesis. *J Dent Res.* 2003 Mar;82(3):162-5. doi: 10.1177/154405910308200303.
7. Mostowska A, Biedziak B, Zadurska M, Dunin-Wilczynska I, Lianeri M, Jagodzinski PP. Nucleotide variants of genes encoding components of the Wnt signalling pathway and the risk of non-syndromic tooth agenesis. *Clin Genet.* 2013 Nov;84(5):429-40. doi: 10.1111/cge.12061.
8. Brook AH. A unifying aetiological explanation for anomalies of human tooth number and size. *Arch Oral Biol.* 1984;29(5):373-8. doi: 10.1016/0003-9969(84)90163-8.
9. Goya HA, Tanaka S, Maeda T, Akimoto Y. An orthopantomographic study of hypodontia in permanent teeth of Japanese pediatric patients. *J Oral Sci.* 2008 Jun;50(2):143-50. doi: 10.2334/josnusd.50.143.
10. Nik-Hussein NN. Hypodontia in the permanent dentition: a study of its prevalence in Malaysian children. *Aust Orthod J.* 1989 Oct;11(2):93-5.
11. Reshitaj A, Krasniqi D, Reshitaj K, Anic Milosevic S. Hypodontia, Gender-Based Differences and its Correlation with other Dental Clinical Features in Kosovar Adolescents. *Acta Stomatol Croat.* 2019 Dec;53(4):347-353. doi: 10.15644/asc53/4/5.
12. Markovic M. *Ortodoncija.* Beograd; 1982.
13. Al Shahrani I, Togoo RA, Al Qarni MA. A review of hypodontia: classification, prevalence, etiology, associated anomalies, clinical implications and treatment options. *World J Dent.* 2013;4(2):117-125.
14. Hall RK. Congenitally missing teeth--a diagnostic feature in many syndromes of the head and neck. *J Int Assoc Dent Child.* 1983 Dec;14(2):69-75.
15. Kokich V. Early management of congenitally missing teeth. *Semin Orthod.* 2005;11:146-151.
16. Kokich VG, Kokich VO. Congenitally missing mandibular second premolars: clinical options. *Am J Orthod Dentofacial Orthop.* 2006 Oct;130(4):437-44. doi: 10.1016/j.ajodo.2006.05.025.
17. Ciarlantini R, Melsen B. Semipermanent replacement of missing maxillary lateral incisors by mini-implant retained pontics: A follow-up study. *Am J Orthod Dentofacial Orthop.* 2017 May;151(5):989-994. doi: 10.1016/j.ajodo.2016.12.012.
18. Garib DG, Alencar BM, Lauris JR, Baccetti T. Agenesis of maxillary lateral incisors and associated dental anomalies. *Am J Orthod Dentofacial Orthop.* 2010 Jun;137(6):732.e1-6; discussion 732-3. doi: 10.1016/j.ajodo.2009.12.024.
19. Rakhshan V. Congenitally missing teeth (hypodontia): A review of the literature concerning the etiology, prevalence, risk factors, patterns and treatment. *Dent Res J (Isfahan).* 2015 Jan-Feb;12(1):1-13. doi: 10.4103/1735-3327.150286.
20. Zilberman Y, Cohen B, Becker A. Familial trends in palatal canines, anomalous lateral incisors, and related phenomena. *Eur J Orthod.* 1990 May;12(2):135-9. doi: 10.1093/ejo/12.2.135.
21. Grahnen H. Hypodontia in the permanent dentition: A clinical and genetical investigation. *Odont Revy.* 1956;7:1-100.
22. Alvesalo L, Portin P. The inheritance pattern of missing, peg-shaped, and strongly mesio-distally reduced upper lateral incisors. *Acta Odontol Scand.* 1969 Dec;27(6):563-75. doi: 10.3109/00016356909026309.

---

\*Corresponding author: Prof. Albena Reshitaj, PhD. Head of the Department of Orthodontics, Medical Faculty School of Dentistry, University of Prishtina, Prishtina, Kosovo. E-mail: albena.reshitaj@uni-pr.edu

23. Koch G, Thesleff I. Developmental disturbances in number and shape of teeth and their treatment. Pediatric Dentistry, 1st ed. Copenhagen: Munksgaard; 2001.
24. Peck S, Peck L, Kataja M. Concomitant occurrence of canine malposition and tooth agenesis: evidence of orofacial genetic fields. *Am J Orthod Dentofacial Orthop*. 2002 Dec;122(6):657-60. doi: 10.1067/mod.2002.129915.
25. Sacerdoti R, Baccetti T. Dentoskeletal features associated with unilateral or bilateral palatal displacement of maxillary canines. *Angle Orthod*. 2004 Dec;74(6):725-32. doi: 10.1043/0003-3219(2004)074<0725:DFAWUO>2.0.CO;2.
26. Baccetti T. Tooth rotation associated with aplasia of nonadjacent teeth. *Angle Orthod*. 1998 Oct;68(5):471-4. doi: 10.1043/0003-3219(1998)068<0471:TRAWAO>2.3.CO;2.
27. Gokkaya B, Kargul B. Prevalence and Pattern of Non-Syndromic Hypodontia in a Group of Turkish Children. *Acta Stomatol Croat*. 2016 Mar;50(1):58-64. doi: 10.15644/asc50/1/8.
28. Badrov J, Gaspar G, Tadin A, Galic T, Govorko DK, Gavic L, Badrov R, Galic I. Prevalence and Characteristics of Congenitally Missing Permanent Teeth among Orthodontic Patients in Southern Croatia. *Acta Stomatol Croat*. 2017 Dec;51(4):290-299. doi: 10.15644/asc51/4/3.
29. Muller TP, Hill IN, Peterson AC, Blayney JR. A survey of congenitally missing permanent teeth. *J Am Dent Assoc*. 1970 Jul;81(1):101-7. doi: 10.14219/jada.archive.1970.0151.
30. Albashaireh ZS, Khader YS. The prevalence and pattern of hypodontia of the permanent teeth and crown size and shape deformity affecting upper lateral incisors in a sample of Jordanian dental patients. *Community Dent Health*. 2006 Dec;23(4):239-43.
31. Maklin M, Dummett CO Jr, Weinberg R. A study of oligodontia in a sample of New Orleans children. *ASDC J Dent Child*. 1979 Nov-Dec;46(6):478-82.
32. Rølling S. Hypodontia of permanent teeth in Danish schoolchildren. *Scand J Dent Res*. 1980 Oct;88(5):365-9. doi: 10.1111/j.1600-0722.1980.tb01240.x.
33. Aasheim B, Ogaard B. Hypodontia in 9-year-old Norwegians related to need of orthodontic treatment. *Scand J Dent Res*. 1993 Oct;101(5):257-60. doi: 10.1111/j.1600-0722.1993.tb01115.x.
34. Arandi NZ, Mustafa S. Maxillary lateral incisor agenesis: a retrospective cross-sectional study. *Saudi Dent J*. 2018 Apr;30(2):155-160. doi: 10.1016/j.sdentj.2017.12.006.
35. Pinho T, Tavares P, Maciel P, Pollmann C. Developmental absence of maxillary lateral incisors in the Portuguese population. *Eur J Orthod*. 2005 Oct;27(5):443-9. doi: 10.1093/ejo/cji060.
36. Celikoglu M, Kazanci F, Miloglu O, Oztek O, Kamak H, Ceylan I. Frequency and characteristics of tooth agenesis among an orthodontic patient population. *Med Oral Patol Oral Cir Bucal*. 2010 Sep 1;15(5):e797-801. doi: 10.4317/medoral.15.e797.
37. Silva Meza R. Radiographic assessment of congenitally missing teeth in orthodontic patients. *Int J Paediatr Dent*. 2003 Mar;13(2):112-6. doi: 10.1046/j.1365-263x.2003.00436.x.
38. Al-Abdallah M, AlHadidi A, Hammad M, Al-Ahmad H, Saleh R. Prevalence and distribution of dental anomalies: a comparison between maxillary and mandibular tooth agenesis. *Am J Orthod Dentofacial Orthop*. 2015 Nov;148(5):793-8. doi: 10.1016/j.ajodo.2015.05.024.
39. Altug-Atac AT, Erdem D. Prevalence and distribution of dental anomalies in orthodontic patients. *Am J Orthod Dentofacial Orthop*. 2007 Apr;131(4):510-4. doi: 10.1016/j.ajodo.2005.06.027.
40. Polder BJ, Van't Hof MA, Van der Linden FP, Kuijpers-Jagtman AM. A meta-analysis of the prevalence of dental agenesis of permanent teeth. *Community Dent Oral Epidemiol*. 2004 Jun;32(3):217-26. doi: 10.1111/j.1600-0528.2004.00158.x.
41. Mani SA, Mohsin WS, John J. Prevalence and patterns of tooth agenesis among Malay children. *Southeast Asian J Trop Med Public Health*. 2014 Mar;45(2):490-8.
42. Stamatou J, Symons AL. Agensis of the permanent lateral incisor: distribution, number and sites. *J Clin Pediatr Dent*. 1991 Summer;15(4):244-6.
43. Coelho ASE, Macho VMP, Andrade DJC, Augusto APC, Areias CMF. Prevalence and distribution of tooth agenesis in a pediatric population: a radiographic study. *Rev Gauch Odontol*. 2012;60:503-8.
44. Delli K, Livas C, Bornstein MM. Lateral incisor agenesis, canine impaction and characteristics of supernumerary teeth in a South European male population. *Eur J Dent*. 2013 Jul;7(3):278-283. doi: 10.4103/1305-7456.115410.
45. Augard, Gayard M. [Statistical study of dental agenesis]. *Ann Odontostomatol (Lyon)*. 1968 Jan-Feb;25(1):21-8. [Article in French].
46. Al-Mulla AA, Mahd TS, Hamid NH. Incidence of hypodontia of permanent teeth. *Tech Res J*. 1990;7:69-80.
47. van den Boogaard MJ, Dorland M, Beemer FA, van Amstel HK. MSX1 mutation is associated with orofacial clefting and tooth agenesis in humans. *Nat Genet*. 2000 Apr;24(4):342-3. doi: 10.1038/74155. Erratum in: *Nat Genet* 2000 May;25(1):125.
48. Magnússon TE. Prevalence of hypodontia and malformations of permanent teeth in Iceland. *Community Dent Oral Epidemiol*. 1977 Jul;5(4):173-8. doi: 10.1111/j.1600-0528.1977.tb01635.x.
49. Pinho T, Maciel P, Pollmann C. Developmental disturbances associated with agenesis of the permanent maxillary lateral incisor. *Br Dent J*. 2009 Dec 19;207(12):E25. doi: 10.1038/sj.bdj.2009.961.
50. Pinho T, Maciel P, Lemos C, Sousa A. Familial aggregation of maxillary lateral incisor agenesis. *J Dent Res*. 2010 Jun;89(6):621-5. doi: 10.1177/0022034510364486.
51. Reshitaj A, Ibishi N, Sejdini M, Berisha N, Mulo XH. Evaluation of the Mesiodistal Crown Sizes of the Remaining Dentition in Patients with Hypodontia, between 12 and 16 Years of Age. *International Journal of Biomedicine*. 2013;3(3):207-210.